


CLAIMS

1. A neurotrophin production/secretion promoting agent which comprises an azole derivative of the formula :



5 wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A represents an acyl group which
 10 may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; X
 15 represents oxygen atom, sulfur atom, or nitrogen atom which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.

2. A neurotrophin production/secretion promoting agent
 20 which comprises a prodrug of an azole derivative or a salt thereof as defined in Claim 1.

3. An agent according to Claim 1, wherein R¹ is a nitrogen-containing heterocyclic group which may
 25 optionally be substituted.

4. An agent according to Claim 1, wherein R¹ is an aromatic heterocyclic group which may optionally be substituted.

30 5. An agent according to Claim 1, wherein R¹ is a nitrogen-containing 5-membered aromatic heterocyclic group which may optionally be substituted.


6. An agent according to Claim 1, wherein R¹ is an imidazolyl group which may optionally be substituted.
- 5 7. An agent according to Claim 1, wherein A is a heterocyclic group which may optionally be substituted, or a hydroxy group which may optionally be substituted.
8. An agent according to Claim 1, wherein A is an aryloxy
10 group which may optionally be substituted.
9. An agent according to Claim 1, wherein A is a phenoxy group substituted with an alkyl group which may optionally be substituted.
- 15 10. An agent according to Claim 1, wherein B is a phenyl group which may optionally be substituted.
11. An agent according to Claim 1, wherein Y is a divalent
20 aliphatic hydrocarbon group.
12. An agent according to Claim 1, wherein X is -O-.
13. An agent according to Claim 1, wherein X is -S-.
- 25 14. An agent according to Claim 1, wherein X is -NR⁴- wherein R⁴ represents a hydrogen atom, a hydrocarbon group which may optionally be substituted, an acyl group which may optionally be substituted, or a heterocyclic group
30 which may optionally be substituted.
15. An agent according to Claim 1, wherein theazole derivative is 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-oxazolepropanol, 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-oxazolebutanol, 4-(4-chlorophenyl)-5-[3-(1-imidazolyl)propyl]-2-(2-methyl-1-

imidazolyl)oxazole, 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-oxazolepentanol, 4-(4-chlorophenyl)-5-[4-(1-imidazolyl)butyl]-2-(2-methyl-1-imidazolyl)oxazole, 3-[3-[4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-oxazolyl]propyl]-1-methyl-2,4-imidazolidinedione, 4-(4-chlorophenyl)-5-[3-(2-methoxyphenoxy)propyl]-2-(2-methyl-1-imidazolyl)oxazole, 4-(4-chlorophenyl)-5-[3-(3-methoxyphenoxy)propyl]-2-(2-methyl-1-imidazolyl)oxazole, 4-(4-chlorophenyl)-5-[3-(4-methoxyphenoxy)propyl]-2-(2-methyl-1-imidazolyl)oxazole, or 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole.

16. An agent according to Claim 1 which is a prophylactic/therapeutic agent for neuropathy.

17. An agent according to Claim 1 which is a prophylactic/therapeutic agent for peripheral neuropathy.

18. A thiazole derivative of the formula :



wherein R^{1a} represents a heterocyclic group which may optionally be substituted; A represents an acyl group which may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.

19. A prodrug of a thiazole derivative or a salt thereof

as defined in Claim 18.

20. A compound according to Claim 18, wherein R^{1a} is a
nitrogen-containing 5-membered aromatic heterocyclic
5 group which may optionally be substituted.

21. A compound according to Claim 18, wherein R^{1a} is an
imidazolyl group which may optionally be substituted.

10 22. A compound according to Claim 18, wherein A is an
aryloxy group which may optionally be substituted.

23. A compound according to Claim 18, wherein B is a phenyl
group which may optionally be substituted.

15

24. A compound according to Claim 18, wherein Y is a
divalent aliphatic hydrocarbon group.

25. A pharmaceutical composition which comprises a
20 thiazole derivative or a salt thereof as defined in Claim
18.


26. A composition according to Claim 25 which is a
neurotrophin production/secretion promoting agent.

25

27. A composition according to Claim 25 which is a
prophylactic/therapeutic agent for neuropathy.

28. A composition according to Claim 25 which is a
30 prophylactic/therapeutic agent for peripheral neuropathy.

29. An oxazole derivative of the formula :



wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A^b represents an aryloxy group which is substituted by an alkyl group and may further be substituted; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.

30. A compound according to Claim 29, wherein A^b is an aryloxy group which is substituted by an alkyl group.

31. A prodrug of an oxazole derivative or a salt thereof as defined in Claim 29.

32. A compound according to Claim 29, wherein R¹ is a nitrogen-containing 5-membered aromatic heterocyclic group which may optionally be substituted.

33. A compound according to Claim 29, wherein R¹ is an imidazolyl group which may optionally be substituted.

34. A compound according to Claim 33, wherein R¹ is an imidazolyl group which may optionally be substituted by a C₁₋₁₀ alkyl.

35. A compound according to Claim 29, wherein B is a phenyl group which may optionally be substituted.

36. A compound according to Claim 35, wherein B is a phenyl

group which may optionally be substituted by halogens.

37. A compound according to Claim 29, wherein Y is a divalent aliphatic hydrocarbon group.

5

38. A compound according to Claim 37, wherein Y is a divalent C₁₋₄ aliphatic hydrocarbon group.

39. A pharmaceutical composition which comprises an
10 oxazole derivative or a salt thereof as defined in Claim 29.

40. A composition according to Claim 39 which is a neurotrophin production/secretion promoting agent.

15

41. A composition according to Claim 39 which is a prophylactic/therapeutic agent for neuropathy.

42. A composition according to Claim 39 which is a
20 prophylactic/therapeutic agent for peripheral neuropathy.

43. 4-(4-Chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole or a salt thereof.

25 44. A crystal of 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole or a salt thereof.

45. 4-(4-Chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(3-methylphenoxy)propyl]oxazole or a salt thereof.
30

46. A crystal of 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(3-methylphenoxy)propyl]oxazole or a salt thereof.


35

47. 5-[3-(4-Chloro-2-methylphenoxy)propyl]-4-(4-

chlorophenyl)-2-(2-methyl-1-imidazolyl)oxazole or a salt thereof.


48. A crystal of 5-[3-(4-chloro-2-methylphenoxy)propyl]-4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)oxazole or a salt thereof.

49. A method for promoting neurotrophin production/secretion in a mammal in need thereof, which comprises administering to said mammal an effective amount of an azole derivative of the formula :




wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A represents an acyl group which may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; X represents oxygen atom, sulfur atom, or nitrogen atom which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.

50. A method for promoting neurotrophin production/secretion in a mammal in need thereof, which comprises administering to said mammal an effective amount of a thiazole derivative of the formula :



wherein R^{1a} represents a heterocyclic group which may optionally be substituted; A represents an acyl group which may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.


51. A method for promoting neurotrophin production/secretion in a mammal in need thereof, which comprises administering to said mammal an effective amount of an oxazole derivative of the formula :



wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A^b represents an aryloxy group which is substituted by an alkyl group and may further be substituted; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.


52. A method for preventing or treating neuropathy in a

mammal in need thereof, which comprises administering to said mammal an effective amount of a thiazole derivative of the formula :



- 5 wherein R^{1a} represents a heterocyclic group which may optionally be substituted; A represents an acyl group which may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may
- 10 optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof.


- 15 53. A method for preventing or treating neuropathy in a mammal in need thereof, which comprises administering to said mammal an effective amount of an oxazole derivative of the formula :



- 20 wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A^b represents an aryloxy group
- 25 which is substituted by an alkyl group and may further be substituted; B represents an aromatic group which may optionally be substituted; and Y represents a divalent

hydrocarbon group or heterocyclic group, or a salt thereof.


54. Use of an azole derivative of the formula :



5 wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; A represents an acyl group which
 10 may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; X
 15 represents oxygen atom, sulfur atom, or nitrogen atom which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof, for the manufacture of a neurotrophin production/secretion promoting agent.

20

55. Use of a thiazole derivative of the formula :




wherein R^{1a} represents a heterocyclic group which may optionally be substituted; A represents an acyl group which
 25 may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, or a carboxyl group which may

optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof,


- 5 for the manufacture of a neurotrophin production/secretion promoting agent.

56. Use of an oxazole derivative of the formula :



- 10 wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; Aᵇ represents an aryloxy group
15 which is substituted by an alkyl group and may further be substituted; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof,
20 for the manufacture of a neurotrophin production/secretion promoting agent.


57. Use of a thiazole derivative of the formula :



- 25 wherein R¹ᵃ represents a heterocyclic group which may optionally be substituted; A represents an acyl group which may optionally be substituted, a heterocyclic group which may optionally be substituted, a hydroxy group which may

optionally be substituted, or a carboxyl group which may optionally be esterified or amidated; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof,
 5 for the manufacture of a pharmaceutical preparation for preventing or treating neuropathy.

58. Use of an oxazole derivative of the formula :



wherein R¹ represents a halogen atom, a heterocyclic group which may optionally be substituted, a hydroxy group which may optionally be substituted, a thiol group which may optionally be substituted, or an amino group which may optionally be substituted; Aᵇ represents an aryloxy group which is substituted by an alkyl group and may further be substituted; B represents an aromatic group which may optionally be substituted; and Y represents a divalent hydrocarbon group or heterocyclic group, or a salt thereof,
 15
 20 for the manufacture of a pharmaceutical preparation for preventing or treating neuropathy.